

## The Effect of Nutritiona Counseling with Booklet Media on Food Waste in Hypertension Patients

Baiq Asmarani Candra Dewi<sup>1</sup>, Luh Suranadi<sup>2</sup>, Made Darawati<sup>3</sup>, Reni Sofyatin<sup>4</sup> and I Gde Narda Widiada<sup>5</sup>

<sup>1-5</sup>Jurusan Gizi Poltekkes Kemenkes Mataram  
<sup>1-5</sup>Jl. Praburankasari Dasan Cermen, Sandubaya Kota  
Mataram Telp./Fax. (0370) 633837  
Email : [renisofyatin@ymail.com](mailto:renisofyatin@ymail.com)

### Article Info

#### Article history:

Received January 2<sup>th</sup>, 2024

Revised February 26<sup>th</sup>, 2024

Accepte March 28<sup>th</sup>, 2024

#### Keyword:

Booklet; Food Waste;  
Hypertension; Low Natrium;  
Nutritiona Counseling;

### ABSTRACT

**Background:** Food waste indicates the success of nutrition services in hospitas; food waste exceeding 20% indicates failure in food management. Nutrition counseling plays a vita role in providing knowledge, attitudes, and behaviors regarding diet so that it can reduce food waste. High Blood Pressure (hypertension) is an increase in blood pressure in the arteries. The number of hypertensive patients at Praya Regiona Hospita in November 2022 will be around 630.

**Objective:** This study's purpose was to determine the effect of nutritiona counseling on changes in plate waste in patients with hypertension at RSUD Praya.

**Research Methods:** This research used a robust quasi-experimenta design, employing a nonrandomized control group pre-and post-test. The sample was carefully selected through purposive sampling, with 22 participants divided equally into treatment and control groups. The data was anayzed using the Mann-Whitney U test. The counseling intervention was delivered through a face-to-face lecture method using booklet media. The study was conducted over one month at Praya Regiona Hospita, ensuring a comprehensive and rigorous approach to the research.

**Research Result:** The study showed differences in patient food waste (staple food, vegetable side dishes, anima side dishes, and vegetables) before and after nutritiona counseling. Still, the difference was insignificant ( $p\text{-vaue} = >0.05$ ). Changes in food waste do not significantly affect staple foods, anima side dishes, vegetable side dishes, and vegetables.

**Conclusion:** Nutritiona counseling using booklet media does not affect food waste in hypertensive patients at Praya Regiona Hospita.

### BACKGROUND

In hospitas, nutrition services are provided and adapted to the patient's condition based on clinica condition, nutritiona status, and body metabolic status. Inpatient nutrition services start with nutritiona assessment, diagnosis, and intervention, including planning, food provision, counseling/education, nutritiona counseling, and nutritiona monitoring and evauation (PGRS, 2013).

The patient's nutritiona condition dramaticaly influences the disease-heaing process. Conversely, the course of the disease can affect the patient's nutritiona condition. The patient's condition often worsens because the body's organs' dietary requirements are unmet. Impaired organ function will worsen with disease and manutrition (Amatsier, 2013).

Efforts to fulfill nutritional needs have been made through nutritional services, such as providing food or diets. The need for nutrients when sick is not only dependent on factors that influence health (age, gender, physical activity, and special conditions) but is also influenced by the severity and type of disease. The need for energy and nutrients also changes during illness and is adjusted to the type and severity of the disease (Amatsier, 2013).

One key strategy to enhance patient food intake is implementing nutritional counseling. The primary goal of nutritional counseling is to boost patient motivation to adhere to the prescribed diet based on their specific health condition. This form of education and counseling can also profoundly impact patient knowledge, which is crucial for effective disease management and therapy adherence.

Nutrition counseling at Praya Regional Hospital is carried out when the patient is going home or, according to a doctor's referral, for a diet consultation for the illness they are suffering from. Counseling at Praya Regional Hospital uses leaflet media adapted to the patient's disease or condition. Besides leaflets, booklets are another medium that can help improve patient knowledge and attitudes. Booklets are also an example of print media that can convey information. A booklet is a small book measuring 14.8x21 cm, with at least five pages but no more than 48 pages outside the cover. The booklet's contents are clear, firm, easy to understand, and enjoyable. Therefore, researchers are interested in conducting counseling using booklet media at Praya Regional Hospital.

High blood pressure (hypertension) is increased blood pressure in the arteries. In general, hypertension is a condition without symptoms, where abnormally high pressure in the arteries causes an increased risk of stroke, aneurysm, heart failure, heart attack, and kidney damage (ASDI 2019). Data obtained in November 2022 at Praya Regional Hospital showed that around 630 patients diagnosed with hypertension were in class 1, 2, and 3 treatment rooms.

Emira's research (2017) showed that there was a change in lunch leftovers in patients with diabetes mellitus at RSI Klaten before and after receiving nutritional counseling, which previously reached 86.21% of leftover staple food, 51.72% of remaining animal side dishes and 41% remaining vegetable side dishes. 38%, and remaining vegetables as much as 44.83%. Then, after receiving nutritional counseling, 34.48% of the remaining staple foods, 10.34% of the remaining animal side dishes, 27.59% of the remaining vegetable side dishes, and 10.34% of the remaining vegetables.

Libriani's research (2020) showed a change in lunch waste in typhoid fever patients before and after receiving nutritional counseling, which previously reached 91.7% of staple food waste, 91.7% of vegetable side dishes, 77.8% of animal side dishes, 97.2% of vegetable side dishes, and fruit 36.1%. Then, after receiving nutritional counseling, the remaining staple foods were 66.7%, vegetable side dishes 75.0%, animal side dishes 52.8%, vegetables 80.6%, and fruit 11.1%.

The data collection results at the Praya Regional Hospital Nutrition Installation showed that inpatients still had food waste. Data on food waste in October 2022 shows that the average food waste in inpatients at Praya Regional Hospital is above the standard the Indonesian Minister of Health set in 2008. The following is data on food waste in October for staple foods: 25%, animal side dishes: 25%, 22.4%, vegetable side dishes: 24.8%, and vegetable food waste: 26.6%.

Based on the above, researchers have researched the effect of changes in food waste before and after nutritional counseling using booklet media for hypertensive patients at Praya Regional Hospital. This research aims to determine the effect of nutritional counseling on food waste in hypertensive patients at Praya Regional Hospital.

## **RESEARCH METHODS**

The research was conducted in class 3 at Praya Hospital in March 2023. It used an experimental method, precisely a quick experiment. The type of Quasy Experiment research method design used is a "Nonrandomized control group pre-and post-test." The sample members in the treatment and control groups were not randomly selected.

The research subjects were patients diagnosed with hypertension who were given a low-sat diet and underwent hospitalization in the class 3 inpatient ward of Praya Regiona Hospita. The tota research sample consisted of 22 people, 11 of whom were in the treatment group and 11 of whom were in the control group. The treatment group was patients diagnosed with hypertension who were hospitalized in Tunjung Room 2, while the control group was patients diagnosed with hypertension who were hospitalized in Tunjung Room 1. In this design, observations were made of the sample before and after being given treatment. The intervention provided in this study was nutritiona counseling twice. In the treatment group, food waste was weighed before and after being given nutritiona counseling using booklet media. In contrast, in the control group, food waste was weighed before and after being given nutritiona counseling without booklet media.

The independent variable in this research is nutritiona counseling, and the dependent variable is food waste. The research instruments used can be seen in the following table:

Data	Instruments	Documentation
Nutrition Counseling	Hypertension Booklet	Photo
Blood pressure	Tensiometer, medica records	Patient biodata form
Leftovers	Food waste recording form	Food waste recording form

## RESULTS AND DISCUSSION

Subjects consisted of 22 patients diagnosed with hypertension who were hospitalized at Praya District Hospita and who met the inclusion criteria, consisting of 11 people in the intervention group and 11 people in the control group. The characteristics of the subjects that have been studied are presented in the following table:

**Table 1 Frequency Distribution of Subject Characteristics**

Karakteristik	Perlakuan		Kontrol		<i>p-value</i>
	n	%	n	%	
<b>Jenis Kelamin</b>					
Laki-Laki	0	0	11	100	0.000
Perempuan	11	100	0	0	
Total	11	100	11	100	
<b>Umur</b>					
30-49	5	45,4	3	27,3	0.429
≥50	6	54,6	8	72,7	
Total	11	100	11	100	

*P-vaue = Mann Whitney test result at  $\alpha = 0.05$*

From Table 1, the Wilcoxon test was carried out in the treatment and control groups to obtain the average waste of staple foods, vegetable side dishes, anima side dishes, and vegetables before and after nutritiona counseling. It was found that the average food waste decreased before and after nutritiona counseling, but a  $p\text{-vaue} > 0.05$  indicates that nutritiona counseling has no effect on staple food waste in hypertensive patients in Tunjung Rooms 1 and 2 at Praya Regiona Hospita.

The results of the Mann-Withney test on staple food leftovers, vegetable side dishes, anima side dishes, and vegetables before and after nutritiona counseling in the treatment group and control group obtained a  $p\text{-vaue} > 0.05$ , which means there was no significant difference in breakfast leftovers in the treatment group and control group in inpatient hypertensive patients at Praya Regiona Hospita.

**Table 2 Distribution of statistica test results of breakfast leftovers in hypertensive patients at Praya Regiona Hospita in the treatment group and control group based on type of food**

Type of Food	Information	Intervention Group (n=11)	Control Group (n=11)	p
		Mean ±SD	Mean ±SD	
Staple food	Before	91,54±87,920	96,15±70,878	0,876
	After	35,69±57,698	53,69±55,182	0,268
	Px	0,38	0,47	
	Δ	55,85	42,46	0,856
Vegetable side dishes	Before	18,00±17,926	14,38±18,035	0,661
	After	5,08±13,048	9,77±15,568	0,367
	Px	0,050	0,176	
	Δ	12,92	4,61	0,689
Anima side dishes	Before	11,38±16,235	8,92±14,829	0,695
	After	2,038±5,0269	1,62±5,824	0,661
	Px	0,075	0,68	
	Δ	9,342	7,3	0,815
Vegetable	Before	21,31±22,321	25,46±22,367	0,871
	After	22,08±21,70	23,85±24,02	0,873
	Px	0,880	0,880	
	Δ	0,77	1,61	0,877

*p*-vaue = Mann Whitney test result at  $\alpha = 0.05$

*px* vaue = Wilcoxon test result at  $\alpha = 0.05$

$\Delta$  vaue = Mann Whitney test result at  $\alpha = 0.05$

From Table 2, the Wilcoxon test was carried out in the treatment and control groups to obtain the average remaining staple food, vegetable side dishes, anima side dishes, and vegetables before and after nutritiona counseling. It was found that the average leftover food decreased before and after nutritiona counseling, but a  $p$ -vaue > 0.05 indicates that nutritiona counseling has no effect on staple food waste in hypertensive patients in Tunjung Rooms 1 and 2 at Praya Regiona Hospita.

The results of the Mann-Withney test on staple food leftovers, vegetable side dishes, anima side dishes, and vegetables before and after nutritiona counseling in the treatment group and control group obtained a  $p$ -vaue > 0.05, which means there was no significant difference in breakfast leftovers in the treatment group and control group in inpatient hypertensive patients at Praya Regiona Hospita.

**Table 3 Distribution of statistica test results for leftover lunch in hypertensive patients at Praya Regiona Hospita in the treatment group and control group based on type of food**

Type of Food	Information	Intervention Group (n=11)	Control Group (n=11)	p
		Mean ±SD	Mean ±SD	
Staple food	Before	87,15±65,239	89,77±72,950	0,938
	After	19,15±26,564	51,54±62,076	0,254
	Px	0,007	0,28	
	Δ	68	38,23	0,290
Vegetable side dishes	Before	20±21,475	18,46±23,635	0,934
	After	3,08±7,610	6,54±16,33	0,870
	Px	0,030	0,128	
	Δ	16,92	11,92	0,790
Anima side dishes	Before	19,46±33,273	5±13,229	0,267
	After	2,69±9,707	17,69±28,412	0,109
	Px	0,138	0,173	
	Δ	16,77	12,69	0,082
Vegetable	Before	49,23±48,760	72,08±56,894	0,393
	After	40,62±47,486	56,85±38,006	0,170
	Px	0,445	0,463	
	Δ	8,63	15,23	0,898

*p*-vaue = Mann Whitney test result at  $\alpha = 0.05$

*px* vaue = Wilcoxon test result at  $\alpha = 0.05$

$\Delta$  vaue = Mann Whitney test result at  $\alpha = 0.05$

From Table 3, the Wilcoxon test was carried out in the treatment group and control group to obtain the average remaining staple food, vegetable side dishes, anima side dishes, and vegetables before and after nutritiona counseling. It was found that the average leftover food decreased before and after nutritiona counseling, but a  $p$ -vaue > 0.05 indicates that nutritiona counseling has no effect on staple food waste in hypertensive patients in Tunjung Rooms 1 and 2 at Praya Regiona Hospita.

The results of the Mann-Withney test on staple food leftovers, vegetable side dishes, anima side dishes, and vegetables before and after nutritiona counseling in the treatment group and control group obtained a  $p$ -vaue > 0.05, which means there is no significant difference in lunch waste in the treatment group and the control group. Control in inpatient hypertension patients at Praya Regiona Hospita.

**Table 4 Distribution of statistica test results of dinner leftovers in hypertensive patients at Praya Regiona Hospita in the treatment group and control group based on type of food**

Type of Food	Information	Intervention Group (n=11)	Control Group (n=11)	p
		Mean ±SD	Mean ±SD	
Staple food	Before	71,77±66,959	91,15±88,169	0,597
	After	48,15±59,23	56,23±50,504	0,619
	Px	0,123	0,202	
	Δ	23,62	34,92	0,856
Vegetable side dishes	Before	15,31±19,202	19,38±16,870	0,552
	After	6,77±12,584	7,23±18,431	0,682
	Px	0,116	0,109	
	Δ	8,54	12,15	0,174
Anima side dishes	Before	9,08±16,142	9,31±18,759	0,793
	After	3,85±13,868	3,69±9,160	0,611
	Px	0,465	0,109	
	Δ	5,23	5,62	0,572
Vegetable	Before	36,38±41,808	50,54±34,508	0,300
	After	36,15±38,583	29,31±20,613	0,979
	Px	0,878	0,091	
	Δ	0,23	21,23	0,105

*p*-vaue = Mann Whitney test result at  $\alpha = 0.05$

*px* vaue = Wilcoxon test result at  $\alpha = 0.05$

$\Delta$  vaue = Mann Whitney test result at  $\alpha = 0.05$

From Table 4, the Wilcoxon test was carried out in the treatment and control groups to obtain the average waste of staple foods, vegetable side dishes, anima side dishes, and vegetables before and after nutritiona counseling. It was found that the average food waste decreased before and after nutritiona counseling, but a *p*-vaue > 0.05 indicates that nutritiona counseling has no effect on staple food waste in hypertensive patients in Tunjung Rooms 1 and 2 at Praya Regiona Hospita.

The results of the Mann-Withney test on staple food leftovers, vegetable side dishes, anima side dishes, and vegetables before and after nutritiona counseling in the treatment group and control group obtained a *p*-vaue > 0.05, which means there was no significant difference in dinner leftovers in the treatment group and control group in inpatient hypertensive patients at Praya Regiona Hospita.

Based on the research results, in terms of age, the 22 subjects in this study were mainly 12 people aged  $\geq 50$  and 10 people aged 30-49 years. One of the factors in the occurrence of hypertension is age; as age increases, the greater the risk of developing hypertension. This is in line with research (Sari et al., 2019), which states that increasing age becomes a risk factor for hypertension. Increasing age causes physica changes in the body, such as thickening of the arteria wals due to the buildup of collagen in the muscle layer, so blood vessels will narrow and become stiff starting at the age of 45. Apart from that, there is also an increase in periphera resistance and sympathetic activity and a lack of baroreceptor sensitivity (blood pressure regulator), and the role of the kidneys, kidney blood flow, and glomerular filtration rate decreases.

Based on the research results, the gender of the 22 subjects in this study was 11 femae and 11 mae. This was because the treatment group in the Tunjung 2 inpatient ward only contained femae patients, and the control group in the Tunjung 1 inpatient ward only contained mae patients. Gender is an interna factor that influences food waste. According to research results (Ronitawati et al., 2020), there is no relationship between gender and patient food waste. In this study, there were 36 respondents,

dominated by 24 women and 12 men. However, in research (Djamauddin, 2018), gender influences the occurrence of food waste, which means there is a relationship.

This research examined staple food waste, animal side dishes, vegetable side dishes, and vegetables. The research results, looking at morning, afternoon, and evening food waste, show that in the treatment group, the average food waste before and after nutritional counseling decreased, as did the control group.

So, based on the results above, even though there was a decrease in food waste before and after nutritional counseling, it is possible that the decrease in food waste was not only influenced by the provision of nutritional counseling; looking at the two sample groups, both showing the same results but with different treatments. Other influencing factors probably caused the decrease in food waste include the patient's preference for certain types of food and menus, the patient's eating habits, consumption of food from outside the hospital, and the patient's inability to finish the food served in the hospital.

Research from Puspita Rahayu (2018) confirms that additional food is usually brought by the patient's family or caregivers outside of the food recommended by the hospital because this food is the patient's favorite. This makes it very likely that the food served to the patient is never finished or not eaten at all.

Food outside the hospital is an external factor; patients often complain about food from the hospital, which is undoubtedly a trigger for food waste; in research (Nafies, 2019), there is a relationship between food outside the hospital and food waste. This is confirmed by the results of research (Ronitawati et al., 2020), which states that food outside the hospital can make patients complete quickly. When mealtime arrives, patients often delay eating so that food quality decreases.

In terms of internal factors, including the patient's psychological state, feeling unhappy and hopeless because of his illness, which can cause loss of appetite (Moehyi, 1992), it can also be caused by the patient's physical condition which determines the type of diet given, if the patient is still weak or there is nausea and vomiting. Hence, it takes a long time to finish the food. The appearance of the food may also cause external factors. For class 3 patients, the food is served with plastic mica; perhaps the method of serving it with plastic mica is less attractive, and the portions are also limited, thus affecting the patient's appetite to finish their food.

Apart from the two factors above, the environment of the treatment place also influences food waste, which is included in the external factors category based on the patient's treatment class, most of which is taken in treatment class III rooms where one room contains eight beds. In treatment Class II, one room contains four beds; in treatment Class I, one room contains two. According to Ronitawati's research results (Ronitawati et al., 2020), there is a relationship between the environment of the treatment area and food waste, where the comfort of the atmosphere around the treatment room can influence the patient's appetite, and this is further strengthened by the results of research (Pontoh et al., 2018)—treatment of patients with leftovers.

Judging from the results of observations and weighing of food waste when compared with treatment and control, both groups experienced a decrease in the average value of food waste, so there was no significant difference between the two groups even though they received different treatment. The statistical test results show no difference in the influence of providing nutritional counseling with booklet media to hypertensive patients at Praya Regional Hospital.

This research is in line with the results of research (Kartini, 2021), which states that when compared with intervention and control food waste, both groups experienced a decrease in the average value of food waste, so there was no significant difference between the two groups even though they received treatment. Different. The results of research on the pre-test intervention and control variables for food waste show that, in general, there is no difference in food waste between the two sample groups. Meanwhile, after the intervention (post-test), it showed no effect of providing nutritional counseling using leaflets on food waste in hypertensive patients.

Likewise, the results of research conducted by (Libriani, 2020) show that the patient's remaining food before counseling consisted of 91.7% of staple foods, 91.7% of vegetable side dishes, 77.8% of animal side dishes, and 97.2% of vegetables and fruit. 36.1%. Meanwhile, the patient's remaining food after counseling was 66.7% staple foods, 75.0% vegetable side dishes, 52.8% animal side dishes, 80.6% vegetables, and 11.1% fruit. There was a change in the patient's remaining food after counseling, but it did not

significantly affect the remaining staple food, vegetable side dishes, animal side dishes, and vegetables. Then, there is a significant influence on the fruit.

Nutrition counseling aims to ensure that clients follow the counselor's suggestions, make positive changes in behavior when solving problems, make decisions, develop awareness, and develop self-acceptance. Counseling can also help clients help themselves by changing their thoughts or lifestyle. Implementing a diet or controlling foods that pose a disease risk still needs to be better understood (Kartini, 2021).

The results of this study do not match the research by Yulianti and Prihatin (2014) regarding differences before and after counseling. The average residual difference was 9.850%, the standard deviation was 5.086, the lowest difference was 7.951%, and the highest was 11.749%, which means there is a difference.

Meanwhile, research by Manik et al. (2019) at RSUD Dr. Soedarso Pontianak stated that providing nutritional counseling using leaflets with leftover food had an effect. The results of the respondents' food waste were obtained by weighing the respondents' food waste starting from food waste, namely rice, animal side dishes, vegetable side dishes, and vegetables. There were more leftover staple foods and vegetables than animal and vegetable side dishes.

More nutritional information will affect the patient's food consumption, so patient eating recommendations are significant. Nutrition counseling is expected to raise patient awareness of food intake. Nutritional counseling can help the disease-healing process by improving nutrition, looking for alternative problem solutions, and choosing the most appropriate way of solving problems for the patient (Kartini, 2021).

The difference in food waste before and after counseling among respondents may be due to respondents still not being focused enough to listen carefully and receive the information that was conveyed during counseling, or it could also be influenced by the condition or location of the counseling, which was carried out in the inpatient room where there were other patients and waiting for patients who are carrying out other activities so that patients find it difficult to understand the information provided (Mappiare, 2006).

This study has limited samples in that the two groups are not the same in terms of characteristics, especially gender, so bias occurs due to the incomparability of the treatment and control groups.

## **CONCLUSIONS**

Based on the results of the research and discussion "The Effect of Nutritional Counseling using Booklets on Leftovers from Hypertension Patients at Praya Regional Hospital," the following conclusions were obtained:

Most research subjects were > over 50 years old, and some were 30-49. The gender of the subjects in the treatment group was all female, and in the control group, all male.

Judging from the research that has been carried out, it shows that in the patient's food waste both before and after nutritional counseling, the most food waste is in the type of vegetable food, namely with an average of 68.9% of food waste.

Nutritional counseling using booklet media does not affect food waste in hypertensive patients at Praya Regional Hospital.

## **REFERENCES**

- Amatsier, Sunita. 2005. *Prinsip Dasar Ilmu Gizi*. Jakarta: Gramedia Pustaka Utama.
- Aritonang, Irianton dan Endah Priharsiwati. 2009. *Manajemen Penyelenggaraan Makanan dan Asuhan Gizi*. Yogyakarta: Leutika Nouvalitera
- Aritonang, Irianton. 2014. *Manajemen Sistem Pelayanan Gizi Swakelola dan Jasaboga di Instalasi Gizi Rumah Sakit*. Yogyakarta: Leutika Nouvalitera



- ASDI. 2019. *Penuntun Diet dan Terapi Gizi Edisi 4*. Jakarta: Buku Kedokteran EGC.
- Djamaluddin, M. (2018). *Analisis Zat Gizi dan Biaya Sisa Makanan pada Pasien dengan Makanan Biasa di RS Sardjito Yogyakarta*. Tesis.
- Departemen Kesehatan RI. 2003. *Pedoman Pelayanan Gizi Rumah Sakit*. Dirjen Bina Kesehatan Masyarakat. Jakarta.
- Departemen Kesehatan RI. 2013. *Buku pedoman Pelayanan Gizi Rumah Sakit, Dirjen Pelayanan Medik, Direktorat Rumah Sakit khusus dan Swasta*. Jakarta.
- Emira. 2017. *Efektivitas Konseling Gizi terhadap Perubahan Sisa Makan Siang pada Pasien Diabetes Mellitus Rawat Inap di RSI Klaten*. Skripsi thesis, Universitas Muhammadiyah Surakarta.
- Iskandar S, Suryani I. 2015. *Pemberian Buku Saku Motivasi Keluarga Dan Pasien Terhadap Penurunan Sisa Makanan Pasien Rumah Sakit*. Gizi Poltekkes Yogyakarta.
- Kartini. (2021). *Pengaruh Konseling Gizi Dengan Menggunakan Media Leaflet pada Pasien Hipertensi Terhadap Tekanan Darah dan Sisa Makanan Pasien Rawat Inap Kelas III Badan Layanan Umum Daerah (BLUD) Rumah Sakit Konawe*. Naskah Publikasi, Politeknik Kesehatan Kendari.
- Kemenkes RI. 2013. *Pedoman Pelayanan Gizi Rumah Sakit*. Jakarta: Kemenkes RI
- Kementerian Kesehatan Republik Indonesia. 2014. *Pedoman Proses Asuhan Gizi Terstandar (PAGT)*. Jakarta: Kementerian Kesehatan Republik Indonesia.
- Kusumaningrum HD, R Yuliah A, Wijaningsih W. 2016. *Pengaruh Konseling Gizi terhadap Pengetahuan Hipertensi Dan Sisa Makanan Pasien Hipertensi Di RSUD DR R Soetijono Blera*. Poltekkes Kemenkes Semarang.
- Libriani Anggeta. 2020. *Pengaruh Konseling Gizi dengan Menggunakan Media Lembar Balik terhadap Perubahan Sisa Makan pada Pasien Demam Tifoid di RSHD Kota Bengkulu*. Skripsi, Poltekkes Kemenkes Bengkulu.
- Lobo EY, Prihatin S. 2014. *Pengaruh Pemberian Konseling Gizi Terhadap Sisamakanan Diet Rendah Garam Di Ruang Rawat Inap Penyakit Dalam Rsud Prof. Dr. W.Z. Johannes Kupang*. Poltekkes Kemenkes Semarang.
- Manik Lisda, Agus Hermansyah, Widyana Lakhsmi Puspita. 2018. *Pengaruh Konseling Gizi menggunakan Media Leaflet terhadap Perubahan Sisa Makanan Pasien Rawat Inap Kelas III di RSUD dr Soedarso Pontianak*. Jurnal, Poltekkes Kemenkes Pontianak.
- Mappiare, A. 2006. *Pengantar Konseling dan Psikoterapi*. Jakarta: Raja Grafindo Persada.
- Menkes RI. 2008. *Keputusan Menteri Kesehatan RI Nomor 129/MENKES/SK/II/2008 tentang Standar Pelayanan Minimal Rumah Sakit*.
- Mourbas I, Novelasari, Nizar M. 2017. *Influence Of Use Of Booklet In Nutritional Concentration On A Change Of Knowledge Patients Diabetes Mellitus In RSUP DR. M. Djamil Padang Year 2017*. Poltekkes Kemenkes Padang
- Nafies, D. (2019). *Hubungan Cita Rasa Makanan dan Konsumsi Makanan dari Luar Rumah Sakit dengan Sisa Makanan Biasa pada Pasien di Rumah Sakit Orthopedi Prof Dr R Soeharso Surakarta*. Skripsi Fakultas Ilmu Kesehatan Universitas Muhammadiyah Surakarta.
- Nuraeni, E. (2019). *Hubungan Usia dan Jenis Kelamin Beresiko dengan Kejadian Hipertensi di Klinik X Kota Tangerang*. Jurnal JKFT Universitas Muhammadiyah Tangerang, 4.
- Notoatmodjo S. 2012. *Promosi Kesehatan dan Perilaku Kesehatan*. Jakarta: Rineka Cipta.
- Persagi, 2010. *Penuntun Konseling Gizi*. Jakarta: PT. Abadi.
- Pontoh, D., Manampiring, A., & Kandou, G. (2018). *Faktor-Faktor Yang Berhubungan Dengan Sisa Makanan Pada Pasien Di Ruang Rawat Inap Rsu Gmim Bethesda Tomohon*. Universitas Sam Ratulangi.

- Puspita, D., & Rahayu, R. (2018). Faktor-faktor yang Berhubungan dengan Perilaku Menyisakan Makanan Pasien Diit DM. *Jurnal Kesehatan Masyarakat Universitas Negeri Semarang*.
- Ronitawati, P., Puspita, M., & Citra, K. (2020). aktor-Faktor Yang Berhubungan Dengan Sisa Makanan Di Rumah Sakit Umum Daerah Koja Jakarta Utara. *Program Studi Gizi, Fakultas Ilmu-Ilmu Kesehatan*.
- Sari, Y. H., Usman, Majid, M., & Sari, R. W. (2019). Faktor-Faktor yang Berpengaruh terhadap Kejadian Hipertensi Pada Lansia di Wilayah Kerja Puskesmas Maiwa Kab.Enrekang. *Jurnal Ilmiah Manusia dan Kesehatan*, 73.
- Supriasa, I D N. 2012. *Pendidikan dan Konsultasi Gizi*. Jakarta. EGC.